

psychological transformations during pregnancy, childbirth, and the first months with the newborn. It is also acknowledged that women can develop mood disorders during this transition to parenthood. The existence of similar deep psychological transformations and the occurrence of mood disorders during men's transition to fatherhood have still to be uncovered.

The Research Program:

The aims of the present fatherhood research program are

- To obtain knowledge of the psychology of men becoming fathers
- To delineate the psychological processes of fathers' relating to their infants
- To develop services appropriate for men becoming fathers

Main Studies:

- Fathers' attachment with their infants
- Mood Disorders in Men Becoming Fathers
- Psychotherapy with men with parental depressions

Sub-studies:

- Fathers and Delivery
- Fathers and Pre-natal Courses
- Grandfathers' Relations with their Sons Becoming Fathers
- Fathers' ability to reflect their infants' states of mind
- Fathers' models of infant caregiving
- Fathers and Biological Fatherhood / Attitudes Towards Donor Insemination

Designs and Participants in the Program:

- Interviews with 72 fathers before and after the birth of the child - about expectations to and experiences with the infant and fatherhood
- Questionnaires from 250 fathers about expectations to and experiences with attending delivery
- Registration of the fathers' presence at 700 deliveries
- Interviews with 10 mothers
- Interviews with 10 midwives
- Questionnaires from 200 fathers about being the biological father
- Questionnaires from 600 fathers about mood disorders
- Psychotherapy with 400 fathers (still accumulating)
- Questionnaires from 210 men undergoing ICSI treatment in three Danish public fertility clinics
- Screening of 3.000 new-born fathers by home nurses in 26 municipalities in 1st to 3rd month post partum
- Screening of 600 fathers to be by GPs in 24th gestation week.
- Analysis of treatment of 47 fathers to be/new-born fathers with perinatal depression.

Head of the research program:

Svend Aage Madsen, Ph.D.

Head of Clinic for Psychology, Play Therapy & Social Counselling
Copenhagen University Hospital, Rigshospitalet, Denmark

Address:

Rigshospitalet, Blegdamsvej 9, DK-2100 Copenhagen - Denmark

Phone no: +45 26212851

E-mail: svendaage@madsen.mail.dk

Co-workers in the research program:

Tina Juhl, Psychologist
Dennis Lind, Clinical Psychologist
Hanne Munck, Associate professor
Marianne Tolstrup, Leading Midwife
Signe Hoffmann Pedersen, Student of Psychology
Frans Svendsen, Psychologist
Inger Poulsen Dutton, Social Counsellor

Helle Dittmann, Play Therapist
Bodil Mortensen, Student of Psychology
Ann Louise Vestergaard, Midwife
Hanne Jensen, Psychologist
Tina Birk Irner, ph.d., Psychologist
Cecilie Skoungaard, Psychologist
Ida Finch-Heidemann, psychologist
Birgitte Bibow, General Practitioner

Summary reports of some projects:

Fathers' Attachment with their Infants

The objective of the main study, *Fathers' Bonds to Their Infants*, is to elucidate fathers' subjective experiences of infant-fatherhood, to analyse the process of bonding from fathers to infants, and thereby to contribute to the theory building on fathers and attachment.

Sample and procedures:

42 randomly selected expectant fathers from the pre-birth ward at the University Hospital of Copenhagen were interviewed three times in two-hour long open-ended interviews. The first interview was conducted 3 months before the birth of the child, the second and third interview were conducted two weeks and five months after the birth of the child.

The results from the interviews were analysed together with transcripts of psychotherapy with 30 fathers who had psychological disturbances about the time of the birth of their infants. These 30 fathers who were treated in psychotherapy were recruited to the study by the departments at the University Hospital, from community health-nurses, and a few fathers made direct contact to the program.

Measures:

The following representations were derived and analysed from the interviews:

- The father's representations of himself as a father
- The father's representations of caregiving from his own childhood (from both his father and his mother)
- The development of the father's representations of the infant before and after birth of the child
- The father's representations of the infant, measured as the father's capability to recognise and respond to his infant's feelings, and needs, as shown in the interview.

Summary of results:

The results from the interviews show a strong correspondence between

- The father's representations of his mother as a caregiver and his capability to recognise and respond to his infant's feelings and needs
- The father's representation of his mother as a caregiver and the representations of himself as a father


There are significant relations ($p < 0,001$) between:

- Fathers having a model of caregiving from their parents characterised by proximity, compassion, and identification
- And fathers expressing a relational fatherhood
- And fathers being able to reflect their infant's states of mind

The same correspondence is not found between the father's representations of the infant and of his own father as a caregiver.

- The father's representations of his infant seem to change through actual interactions and experiences with his infant: Most of the fathers' representations of the child before the birth are representations of a pre-school child, who is capable of verbal contact and ball-playing. The representations change for almost all the fathers after the birth to representations of the actual infant. A small group of fathers' representations do not change - they remain representations of the child as a pre-schooler. This group of fathers shows difficulties in recognising and responding to their infant's feelings and needs.

These results underline the importance of focusing on the father's representations of his mother as a caregiver in understanding the bonding process between fathers and their infants. It also stresses the importance of early interactions between father and newborn to establish the development of the father-infant-relationship in the father-bonding perspective.



Fathers and Delivery

The sub-study on fathers and delivery was conducted in Denmark in 1998-99. Research results from this study are presented in summary below:


The expectations, thoughts, wishes, fears and actual experiences of the fathers as well as the midwives' experiences of fathers attending the delivery process were investigated.

Methods and participants:

Consecutive registration in a three-month period (2.800 deliveries a year) of the fathers' presence at delivery. In all 165 fathers completed questionnaires in the 38th week of the women's pregnancy and again following the delivery. A semi-structured interview was given to 23 fathers before and after the delivery. A fifth of the midwives randomly selected were given a semi-structured interview.

Results:

- The fathers were present in 88.1% of all deliveries (n=698). The reasons for not participating seem related to either an upbringing in a Third World country (6%), special circumstances at the delivery, or the woman being single (3 %). Subtraction of these aspects leads to a father participation rate of 97%.
- The fathers stated that they wanted to be present for their own sake and not only to support their partners, and that they were satisfied to have been there (98%), and felt that they had played an important role.
- 75% of the fathers participated in pre-natal courses and 88% of the fathers participated in prophylactic consultations. Almost all of them found it beneficial. Two-thirds felt personally welcome, but 54% did not feel directly invited and 40% did not experience that the midwives directly addressed them during the consultations.



Discussion:

Our findings thus show a strong involvement from the fathers during pregnancy and delivery. At the same time our findings point to an apparent discrepancy between the attitudes of the official health authorities (The Danish Health Board) and the families, concerning the father's role and the importance of his participation in the birth of his child.

The official health authorities hardly mention the father in their publications and general directives. Likewise, the traditional hospital routines do not allow the father to stay at the hospital overnight after delivery with the mother and their newborn infant in contrast to the wish of the fathers and their wives. Such a policy indirectly supports the view of the fathers' presence as being of little importance, ignoring the involvement of the fathers, which is clearly shown in our study.

Mood Disorders in Men Becoming Fathers

Fathers and Postnatal Depression:

Men's Psychological Transition into Fatherhood

Overall Findings:

The study, which involved 600 men, indicates that around **7 percent** of fathers suffer from postnatal depression.

This is the equivalent of around **225-250 fathers** of babies born at *Rigshospitalet* who are suffering from postnatal depression on a yearly basis. Around 3.400 births take place at *Rigshospitalet* annually.

The current annual no. of births in Denmark is around 65.000. Hence on a national level the numbers indicate that roughly **4-4.500 fathers** suffer from postnatal depression.

About the study:

The study used two different scales to detect depression: 1) the 'Edinburgh Postnatal Depression Scale' (EPDS), traditionally used to screen for postnatal depression, and 2) the 'Gotland Male Depression Scale' (GMDS), normally used to assess so-called masculine states of depression.

The two scales list the following conditions as possible indicators of depression:

EPDS ('Traditional' depression)	Gotlandskala ('Male' depression)
<ul style="list-style-type: none"> • Unable to laugh or be humorous • Unable to look forward to things • Blaming myself unnecessarily when things go wrong • Feeling anxious or worried for no good reason • Feeling scared or panicky for no good reason • Things get on top of me • Have been so unhappy that I have had difficulty sleeping • Feeling sad or miserable • Feeling so unhappy that I have been crying • The thought of harming myself has occurred to me 	<ul style="list-style-type: none"> • Lowered stress tolerance/feeling more stressed • Feeling aggressive/outward-reacting/low impulse control • Feeling burned out and empty • Inexplicable fatigue • Feeling irritable, restless and frustrated • Difficulty making everyday decisions • Sleeping too much/too little/sleeping fitfully • Difficulty falling asleep/early waking • Feeling agitated/anxious/uncomfortable • Excessive use of pills and/or alcohol • Excessive activity/working hard, incessantly/ jogging etc. • Altered behaviour makes me difficult to be around • Feeling that you perceive yourself/others perceive you as gloomy, negative • Feeling that you perceive yourself/ others perceive you as moaning, self-pitying

The 600 men who participated in the study were recruited through antenatal classes and midwife consultations at *Rigshospitalet*. The response rate among the men who received the questionnaire was 91%.

The study was carried out from February to December of 2005 and included 600 fathers whose child was born at *Rigshospitalet*. The men's condition was assessed by asking them to fill out a questionnaire roughly 6 weeks after the birth of their child.

Up to this point the *Fædreforskningsprogrammet* - the Paternal Research Programme - at *Rigshospitalet* has received about **70 fathers** for psychotherapeutic treatment for postnatal depression. In the course of this work it has become apparent that postnatal depressions may arise both before the birth and for a significant period afterwards.

The project receives funding from the EU-Commission under '*Community Framework Strategy on Gender Equality 2001-2005*'. The Project has been carried out by a research group consisting of Chief Psychologist *Svend Aage Madsen, Ph.D.*, Psychologist *Tina Juhl* and Midwife *Ann Louise Vestergaard* - all members of *Rigshospitalet's* Psychological and Psychosocial Research Unit.

Further Details:

When we used the questionnaire traditionally used to identify postnatal depression in women - EPDS - it was estimated that 5,0 % of men examined suffer from postnatal depression. Using a specific scale for 'Male depression' - the Gotland Scale - an estimated 3,4 % of men were found to be suffering from postnatal depression.

Making use of both scales the results show roughly 7% of fathers to be suffering from postnatal depression (the fathers who register as depressive according to both scales are counted only once). The number includes both first-time fathers and some who have previously had children.

The numbers show the following distribution amongst the 544 fathers who responded:

EPDS	27 (5,0% ¹)
Gotland	18 (3,4% ²)
Postnatal Depression including EPDS + Gotland	34 (6,5%³)
Valid answers: ¹ 537, ² 524, ³ 520	

In comparison, international research shows that 10% of women suffer from postnatal depression as registered by the EPDS about 6 weeks after the birth.

Comparing postnatal depression with other forms of depression, recent Danish research shows that 3,5% of 20 to 50-year-old men suffer from depression of moderate to medium severity at any given time. Twice as many women as men are diagnosed as suffering from some form of depression.

There are a number of indicators that men's psychiatric problems are generally under-diagnosed and that they hence receive insufficient treatment in this area.

Perspectives:

The fact that men do suffer from postnatal depression was demonstrated from the results of the research project 'Fathers' Bonding with their Infants' (2) and was first publicly announced in Denmark in 2002. Since then work has been going on at *Rigshospitalet* to investigate this phenomenon, including i.a. psychological treatment of men suffering from this condition.

The current survey indicates the necessity of training health-care staff -who come into contact with families during pregnancy, birth and infancy - to also be aware of the psychological state of fathers and to identify those suffering from postnatal depression.

Furthermore the method utilized here to identify postnatal depression in men will also be useful in developing tools to improve general diagnosis of depression in men.

It appears that for now *Rigshospitalet* is the only place in Denmark offering specific treatment – and the therapy we have been able to offer here at *Rigshospitalet* is underwritten by EU project-funds for a limited period of time only. Thus the survey also implies a need to procure treatment for the roughly 4-4.500 men a year who are estimated to suffer from this condition nationwide

This finally indicates a need for training treatment providers for men suffering from postnatal depression



(1) Olsen, L., Mortensen, E. & Bech, P. (2004) *Prevalence of major depression and stress indicators in the Danish general population*. Acta Psychiatrica Scandinavica: 109: 96-103.

(2) Madsen, S.Aa., Lind, D. & Munck, H. (2002) *Fædres tilknytning til spædbørn*. Copenhagen: Hans Reitzel Publishers.

Psychological aspects of male infertility in a Scandinavian ICSI population

A.T. Mikkelsen, S.Aa. Madsen, L. Ohrt, L. Dessing & P. Humaidan.

Introduction:

The interest in male reproduction and psychological reactions related to ART has increased during recent years, coinciding with the use of intracytoplasmic sperm injection (ICSI). Previous studies have shown that male infertility may cause a loss of self-esteem and impaired gender identity. Moreover, male patients are inclined to ignore their own needs in order to show consideration for the partner, as focus will invariably be on the female partner. The aim of this study was to document the experiences of infertile Danish men as individuals, partners and patients, respectively, when ICSI treatment was the only possible way to establish fatherhood. Furthermore, to identify the possible need for additional psychological communicative support of the infertile man in the care process.

Materials and methods:

A self-report questionnaire with structured and open-ended questions was completed by 210 men, undergoing ICSI treatment in three Danish public fertility clinics. The questionnaire consisted of 69 questions in total, including three issues: individual perception of reduced sperm quality, marital consequences and communication with health professionals in the clinic. All questionnaires were completed in private in the clinic, while the female partner rested after the oocyte retrieval.



Results:

Among the 210 participants, 95.7% found fatherhood to be important, and the need for ICSI was not important for the perception of fatherhood. 41% claimed that the need for fertility treatment did not negatively affect the relationship with their partner; on the contrary 51% found that it had strengthened the relationship. Interestingly, in 37% of the participants the reduced sperm quality affected their perception of masculinity. Participants were generally satisfied with the care, and only 21% expressed concerns about ICSI. However, 62% found that there was a need for a deeper dialogue with the staff concerning the experience regarding reduced sperm quality, and 72% lacked information regarding psychological consequences of male infertility in general.

Participants wished health professionals to show openness in questions concerning reactions to infertility, infertility treatment and the role of the male partner.

Conclusions:

Infertile men wish to be met by health professionals on equal terms with their female partner. The ICSI procedure per se is of no importance for the perception of fatherhood. Interestingly, the reduced sperm quality affected one third of the participants negatively in the perception of their masculinity. Our data suggest that infertile Danish men are prepared to a greater degree of openness and information regarding male infertility and the relating psychological consequences.

Authors:

A.T. Mikkelsen¹, S.A. Madsen², L. Ohrt³, L. Dessing⁴, P. Humaidan¹.

¹Skive Regional Hospital, Fertility Clinic, Skive, Denmark.

²Copenhagen University Hospital Rigshospitalet, Ward 4074, Copenhagen, Denmark.

³Braedstrup Regional Hospital, Fertility Clinic, Braedstrup, Denmark.

⁴Holbaek Regional Hospital, Fertility Clinic, Holbaek, Denmark.

Themes in psychotherapy with men with parental depressions

Up to now – end of 2016 – 400 men with parental depressions have been treated in our clinic.

400 men with parental depression in psychotherapy

Time of referral	%
During pregnancy	40
Around delivery & confinement (< 1 month)	10
Infancy > 1 month	50



From the therapeutic work and with the background results from the Fædreforskningsprogrammet (Fatherhood Research Program) (Madsen, Lind & Munck 2002; Madsen 2009), we have found that the fathers are characterized by meeting up with the conditions, issues and reactions listed below:

- Anger - the man's extroverted, negative states of mind often find expression in aggression, sometimes against himself
 - Withdrawal - often withdrawal from close relations is seen as a way to relieve pain
 - Past and present relations with caregivers - the man often feels uncared for
 - Relations with the his infant - often the aggressions are directed towards the child
 - Relations with spouse - sometimes the aggressions are directed towards the wife
 - Ambivalence between longing for closeness and attachment on the one hand and autonomy and detachment on the other
- Images of masculinity related with the infant's need for care.

The method used is called 'Minding the parent-child-relationship'.

Treatment modalities:

A two-pronged psychotherapy working with

- the man's past and present relations with caregivers
- alternating with his relations with the infant

Early detection at the GPs of perinatal depression in mothers and fathers to be

Background

The Danish Board of Health states (2013): "The incidence of postpartum depression has been found to be about 10% in mothers up to 6 months after birth and about 6% in the fathers". Men and women's pre- and postnatal depression may have impact on a child's social-emotional development. There is a need to detect and treat the depression as early as possible to avoid negative effects on the child.

Objectives

The objectives of the project were to investigate whether

- perinatal depressions are detectable during pregnancy
- it is possible to identify fathers with perinatal depression,
- it is possible to conduct a screening at the GP's in the 24th week of pregnancy,
- the mothers and fathers to be, who score above the cut off, want to be offered psychotherapy. Finally, it was the aim to compare figures and symptom profiles from this study with results from screening of mothers and fathers 6 to 8 weeks after the birth of their child.

Finally, it was the aim to compare prevalence and symptoms of perinatal depression among parents to be with parents 6 to 8 weeks after the birth of their child.

Material/Methods

30 practitioners in the City of Copenhagen have in a 1½ year period screened 1,200 expectant mothers and fathers for perinatal depression in the women's 24th week of pregnancy. The screening instrument for both parents were a combination of the Edinburg Post-Natal Depression Scale (Cox et. al., 2003) and the Gotland Scale of Male Depression (Rutz et. al., 2002) with selected questions from The Masculine Depression Scale (Magovcevic et al., 2009) and the Father Attachment Interview (Lind et. al., Unpublished manual)



All responders who scored above the cut off were referred to psychological treatment before the child's birth. At the same time, statistical data were collected for exploration of the occurrence of depression, symptom profiles and gender differences.

Results and Conclusion

In 1204 screenings in the 24th week of pregnancy a total of 9.8% scored above cut off, i.e. 11.1 % of women and 8.0 % of men. The project is unique internationally in examining whether it is possible in general practice to detect signs of postpartum depression during pregnancy and in screening fathers to be for perinatal depression. We conclude that it is possible to detect postpartum depression already during pregnancy. After completing analyses, it is expected to form a basis for introducing screening parents to be for perinatal depression in general practice.

Litterature

Cameron, E., Sedov, I. & Tomfor-Madsen, L. (2016) Prevalence of paternal depression in pregnancy and the postpartum: An updated meta-analysis'. *Jrn Affective Disorders*. <http://dx.doi.org/10.1016/j.jad.2016.07.044>

Cox, J. & Holden, J. (2003): *Perinatal Mental Health: A Guide to the Edinburgh Post-natal Depression Scale*. London: Gaskell.

Danish Board of Health, The (2013) *Anbefalinger for svangreomsorgen [Guidelines for Perinatal Care]*. Copenhagen.

Lind, D., Madsen, S. Aa., & Munck, H. (1998): *Father attachment interview*. Unpublished manual. Copenhagen University Hospital, Rigshospitalet & University of Copenhagen Fædreforskningsprogram.

Madsen, S.Aa. & Juhl, T. (2007). Paternal depression in the postnatal period assessed with traditional and male depression scales. *Int Journ Men's Health & Gender*. 4, I, 26-31.

Madsen, S.Aa. (2008) Spædbarnet i faderens verden. *Psyke & Logos*, 7, 2, 669-687.

Magovcevic, M. & Addis M. (2009): The Masculine Depression Scale: Development and Psychometric Evaluation.

Paulson J., Dauber S. & Leiferman J. (2006): Individual and Combined Effects of Postpartum Depression in Mothers and Fathers on Parenting Behaviour. *Paediatrics*, 118, 659-668.

Rutz, W., Rihmer, Z. & Dalteg, A. (2002): The Gotland Scale for assessing male depression. *Nord J Psychiatry*. 56(4): 265-71.

Authors

Svend Aage Madsen, Birgitte Bibow, Anette Sonne Nielsen, Sarah Tranekær Rasmussen, Hans Christian Møller,

Index List: Prevention of mental problems, parenting, infants, postnatal depression

Developing the 'Minding the father-child-relationship' model for treatment of fathers with perinatal depressions.

Around 10 pct. fathers suffer from perinatal depression. This disorder should be viewed as a relational disturbance rooted in relations from the father's own upbringing, constituting itself in relations as a parenthood disturbance also influencing the mother and influences relations in future, namely by

having impact on the child's development. Psychotherapy therefore must approach this suffering as a relational disturbance.

The treatment model builds upon the Cochrane analysis (Dennis & Hodnett 2007) concluding: "Overall, psychosocial and psychological interventions are effective treatments for postpartum depression." The treatment model is developed through a modified 'Mentalising-based psychotherapy' (Allen, Fonagy & Bateman 2008), and inspired by 'Minding the Baby' (Sadler, Slade & Meyers 2006).

The treatment is tested for effects on the men's symptoms and on the father-child-attachment.

The treatment model is described in the publication: Madsen, S.Aa. (2018) *Guide til behandling af fædre med fødselsdepression [Guide for Treatment of Fathers with Perinatal Depression]*. København: Rigshospitalets Fædre-forskningsprogram.



The treatment model engages with 'male specific' mental states, because many men in their upbringing - and in the cultural and societal expectations of masculinity - are confronted with and have psychologically internalised an ambivalence between autonomy and freedom on the one hand, and closeness and attachment on the other. This ambivalence becomes sharpened in men's mental states during the process of being a parent, and it is a vital issue in postnatal depression in men. This ambivalence is often displayed in a man's tendency to withdraw himself from relationships in the hope that he will find relief in distancing himself from close family. It also appears in many men's wanting to quickly get away from agony, sometimes by alcohol abuse, irrational actions, etc. Finally, this also appears in some men's tendencies to act out in angry out-bursts, violence, or different abuse.

Besides the mainstream psychotherapy that builds upon psychodynamic, mentalising, narrative, or cognitive methods it is therefore useful to include the above-mentioned topics in a treatment model suited for men with perinatal depression and men's other family related disorders.

Publications from the research program (selected):

- Madsen, S.Aa. (1995). *Forskning i fædres tilknytning til deres spædbørn*. Paper præsenteret på Høstseminar i Nordiska Föreningen För Spädbarns Utveckling på Københavns Universitet d. 22.-23. september
- Madsen, S.Aa. (1996). Fædre, fødsler og psykiske reaktioner. *Tidsskrift for Jordemødre* (5), 'Temnummer om mænd', 30-33.
- Madsen, S.Aa., Rode, U. & Sørensen, L. (1998) *FAR - En folder til mænd på vej til at blive fædre*. Allerød: SCA Mölnlycke.
- Madsen, S.Aa. (1998). "Forskning i fædres tilknytning til deres spædbørn". Paper præsenteret på *Årskonference i Nordisk Forening for Spædbarnsudvikling* på Rigshospitalet i København.
- Madsen, S.Aa. & Munck, H. (1999). Fædres deltagelse i fødsler – nærvær eller afstand. In: Ervø, Hove & Nik-Meyer (Eds.) *Mænd på vej*. København: Ligestillingsrådet.
- Madsen, S.Aa. , Munck, H. & Tolstrup, M. (1999). *Fædre og fødsler*. København: Frydenlund Grafisk.
- Madsen, S.Aa. , Munck, H. & Tolstrup, M. (1999). Fædres tilstedeværelse ved fødsler. *Tidsskrift for Jordemødre*, 8, 15-19.
- Madsen, S.Aa., Munck, H. & Lind, D. (2000). Faderskab og tilknytning – en oversigt over tilknytningsforskningen i et fædreperspektiv. In: Elsass & Lunn *Psykologi i forvandling – et jubilæumsskrift fra Institut for Klinisk Psykologi* København: Københavns Universitet
- Munck, H., Lind, D. & Madsen, S.Aa. (2000). Fædre og tidlig intervention. In: Elsass & Lunn. *Psykologi i forvandling – et jubilæumsskrift fra Institut for Klinisk Psykologi* København: Københavns Universitet
- Madsen, S.Aa. (2000) Far og barn i fokus. *Tidsskrift for Jordemødre* (1), 5.
- Madsen, S.Aa. (2000). Man kan hvis man letter røven. In: Mauritson, M. & Mauritsen, M. *Far for første gang*. København: Aschehoug.
- Lind, D, Madsen, S.Aa. & Munck (2000). Fathers and Delivery. *Infant Mental Health Journal* (21), 4-5, 241.
- Lind, D., Madsen, S.Aa. & Munck, H.(2000). Fædeforskning: Fra rolletænkning til dynamisk teori. In: Elsass & Lunn. *Psykologi i forvandling – et jubilæumsskrift fra Institut for Klinisk Psykologi* København: Københavns Universitet
- Madsen, S.Aa., Munck, H., Pedersen, S. & Tolstrup M. (2000). Fædre og fødselsforberedelse. *Tidsskrift for Jordemødre*,(110), 8, 4-9.
- Munck, H., Madsen, S.Aa. & Lind, D. (2000). *Fathers' relations to their infants*. Presented at WAIHM's 7th World Congress in Montreal, Canada.
- Pedersen, S. Hoffmann. (2000) *Jeg skal være far – om fædres forberedelse til fødslen og livet med et spædbarn* København: Forskningsprogrammet 'Fædres relationer til deres spæd- og småbørn'.
- Madsen S.Aa. & Munck, H. (2001). Une étude au Danemark. La présence des pères à l'accouchement. *Santé Mentale au Québec*, XXVI.(1), 27-38.
- Madsen, S. Aa.. (2001) Fædre og spædbørn. In: Andersen, H. (Ed.) *Aschehougs store babybog*. København: Aschehoug.

- Madsen, S. Aa., Lind, D & Munck, H. (2002). *Fædres tilknytning til spædbørn*. København: Hans Reitzels Forlag.
- Madsen, S.Aa. & Munck, H. (2002). *Men as fathers present at deliveries in hospital*. Book of Abstracts, 2 nd World Congress on Men's Health: Sex & Gender Matter From Boys to Men – From Science to Practice, October 25 - 27, 2002, University of Vienna, Vienna, Austria
- Munck, H., Madsen, S.Aa. & Lind, D. *Fathers' thoughts and feelings towards becoming and being a father*. Paper presented at WAIHM 8th World Congress in Amsterdam, Holland, July 2002.
- Madsen, S. Aa.. (2003) Hvor er barnets far? Om forandringer i spædbarnsfaderskabet. In: Baagøe-Nielsen, S. & Hjort, K. (red.) *Mænd og Omsorg*. København: Hans Reitzels forlag.
- Madsen, S.Aa. (2003) Svangreomsorgen og barnets far. Statusartikel. *Ugeskrift for Læger*, vol. 165, 46., 4423-5.
- Madsen, S.Aa. (2003) Fædre i svangreomsorgen. *Månedsskrift for praktisk lægegering*, Vol. 81, 9: 1229-1234.
- Madsen, S.Aa. & Lind, D. (2003) Men's mood disorders in connection with the birth of their child. Oral and Poster presentation at The 3rd World Congress on Men's Health, Vienna October 24th-26th.
- Lind, D. & Madsen, S.Aa. (2003) Men in Psychotherapy. Poster presentation at The 3rd World Congress on Men's Health, Vienna October 24th-26th.
- Madsen, S.Aa. (2003) Fædrene i tilknytningsforskningen. *Temanr. Om tilknytning. Psyke & Logos*, 2.
- Madsen, S.Aa. & Lind, D. (2004) The Infant in the Father's World. Symposium lecture at World Association for Infant Mental Health. – World Congress, Melbourne, Australia – January 14-17
- Barnes J, Sutcliffe AG, Kristoffersen I, Loft A, Wennerholm U, Tarlatzis BC, Kantaris X, Nekkebroeck J, Hagberg BS, Madsen S.Aa., and Bonduelle M (2004) The influence of assisted reproduction on family functioning and children's socio-emotional development: Results from a European study. *Human Reproduction*. Jun 2004; 19: 1480 - 1487.
- Madsen, S.Aa. (2005) *A New Fatherhood Emerging Fathers' Relations With Their Infant*. Lecture at School of Social Studies, Masaryk University in Brno, Czech Republic, November 2004.
- Madsen S.Aa. (2005). *Male sensitive communication on illness, lifestyle, and being a parent in a hospital setting*. World Congress on Men's Health and Gender. 1st October, Vienna, Austria.
- Madsen, S.Aa. (2004) Fædre og barselsorlov. *Kvinden og Samfundet* 120, 1. S. 4-5
- Madsen, S.Aa. (2005) Kan mænd få fødselsdepressioner? 'Spørg Riget', *Politiken* 11. sep. 12-13.
- The Web-site: <http://www.european-fatherhood.com/>
- Madsen, S. & Juhl, t. (2006) Mænd kan også få fødselsdepressioner. *Psykologisk Set*, 23, vol 62, 4-11.
- Madsen, S.Aa. & Juhl, T. (2007). Paternal depression in the postnatal period assessed with traditional and male depression scales. *Int Journ Men's Health & Gender*. Vol 4 Issue I, March, 26-31.

- Madsen, S.Aa., Lind, D. & Munck, H. (2007). Men's Abilities to Reflect Their Infants' States of Mind - Interviews with 41 new fathers on experiences of parenthood. *Nordic Psychology*, 59,(2), 149-163.
- Madsen, S.Aa. (2007) Fædre med fødselsdepression. *Forældre og fødsel. Tema-nummer om efterfødselsreaktioner. Nr. 5, September. 35-38.*
- Madsen, S.Aa. (2008) "Psychische Belastungsstörungen bei Männern, die Väter werden - Psychodynamik, Häufigkeiten und Auswirkungen auf die Praxis. In: Schäfer, E., Abou-Dakn, M. & Wöckel, A. (eds.): (Eds.) Vater werden ist nicht schwer? Zur neuen Rolle des Vaters rund um die Geburt. Edition Psychosozial. Berlin: Psycho-sozial Verlag.
- Madsen, S.Aa. (2008) Spædbarnet i faderens verden. Temanr. Om spædbarnspsykologi. *Psyke & Logos*, 7, 2, 669-687.
- Madsen, S.Aa. (2009) Men's Mental Health: Fatherhood and Psychotherapy. *The Journal of Men's Studies. Vol 17, 1 (winter)*, 15-30.
- Madsen, S.Aa. & Burgess, A. (2010) Becoming a father – post natal depression in men. In: White, A. & Conrad, D. (eds.) *Promoting Men's Mental Health*. London: Radcliffe Publishing.
- Madsen, S.Aa. (2010) Between autonomy and attachment. Psychotherapy for men with postnatal depression. In: Blazina, C & Miller, D. (eds). *An International Psychology of Men. Theoretical Advances, Case Studies, and Clinical Innovations*. New York: Routledge. (in press).
- Madsen, S.Aa. (2010). *Treatment of Anger in New Fathers and Fathers to Be*. Article from Daphne III –program project Violence Intervention in Specialist Healthcare (VISH) granted by the European Commission
- Madsen, S.Aa. & Mikkelsen, A. (2010). *Mænd går også i fertilitetsbehandling*. København: MSD.
- Mikkelsen, A., Madsen, S. A. & Humaidan, P. (2012) Psychological aspects of male fertility treatment. *Journal of Advanced Nursing Article first published online: 10 DEC 2012. DOI: 10.1111/jan.12058*
- Ammizbøll, J., Gullestrup, L., Lindved, K., Madsen, S.Aa., Videbech, P. & Vinter, M. (2012) *Fødselsdepression – der er hjælp at få!*. København: Frydenlund.
- Madsen, S.Aa. (2014) *Mænds sundhed og sygdomme*. København: Samfundslitteratur.
- Madsen, S.Aa. (2014) Psykologiske aspekter ved graviditet, fødsel og barsel. I: Uldbjerg, N., Damm, P. & Jørgensen, J. *Obstetrik – en grundbog*. København: Munksgaard.
- Madsen, S.Aa. (2015) Men as patients. *Trends in Urology and Men's Health* 6, 3 May/June, 22–26.
- Madsen, S.Aa. (2017) Opsporing af fødselsdepression før fødslen. *Sundhedsplejersken*, 3, 39, s. 32-35.
- Madsen, S. Aa. & Prentow, T. (2018) Fædrene og sundhedsplejersken – og sundhedsplejersken og fædrene. *Sundhedsplejersken*, 01,40, 14-17.
- Madsen, S.Aa., Bibow, B. & Eliasson, S. (2018) Almen praksis kan opdage fødselsdepression allerede under graviditet. *Månedsskrift for almen praksis*, maj, 424-431

Madsen, S.Aa. (2018) *Guide til behandling af fædre med fødselsdepression*. København: Rigshospitalets Fædreforskningsprogram.

Mikkelsen, A., Madsen, S. A. & Humaidan, P. (2019) Den infertile mand skal ind fra sidelinjen. *Politikens Kroniken* 23. september. Politiken.

Prentow, T.S. & Madsen, S.Aa (2020) Guide til bedre inddragelse af fædre i sundhedsplejen. *Sundhedsplejersken*, 2, april, 20-24.

Madsen, S.Aa (2020) Forældre sammen – en brugsbog i svangreomsorgen. *Månedsskrift for Almen Praksis*, 4, april, 25-31.

Prentow, T. & Madsen, S.Aa. (2021) *Far for livet – En bog til manden som far*. København: Muusmanns Forlag.